

Marginal and conditional distributions of multivariate normal distribution

Assume an n -dimensional random vector

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix}$$

has a normal distribution $N(\mathbf{x}, \mu, \Sigma)$ with

$$\mu = \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} \quad \text{and} \quad \Sigma = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}$$

where \mathbf{x}_1 and \mathbf{x}_2 are two subvectors of respective dimensions p and q with $p + q = n$. Note that $\Sigma = \Sigma^T$, and $\Sigma_{21} = \Sigma_{12}^T$.

Theorem 4:

Part a The marginal distributions of \mathbf{x}_1 and \mathbf{x}_2 are also normal with mean vector μ_i and covariance matrix Σ_{ii} ($i = 1, 2$), respectively.

Part b The conditional distribution of \mathbf{x}_i given \mathbf{x}_j is also normal with mean vector

$$\mu_{i|j} = \mu_i + \Sigma_{ij}\Sigma_{jj}^{-1}(\mathbf{x}_j - \mu_j)$$

and covariance matrix

$$\Sigma_{i|j} = \Sigma_{ii} - \Sigma_{ij}\Sigma_{jj}^{-1}\Sigma_{ji}$$

Proof: The joint density of \mathbf{x} is:

$$f(\mathbf{x}) = f(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{(2\pi)^{n/2}|\Sigma|^{1/2}} \exp\left[-\frac{1}{2}(\mathbf{x}-\mu)^T\Sigma^{-1}(\mathbf{x}-\mu)\right] = \frac{1}{(2\pi)^{n/2}|\Sigma|^{1/2}} \exp\left[-\frac{1}{2}Q(\mathbf{x}_1, \mathbf{x}_2)\right]$$

where Q is defined as

$$Q(\mathbf{x}_1, \mathbf{x}_2) = (\mathbf{x} - \mu)^T \Sigma^{-1} (\mathbf{x} - \mu)$$

$$\begin{aligned}
&= [(\mathbf{x}_1 - \mu_1)^T, (\mathbf{x}_2 - \mu_2)^T] \begin{bmatrix} \Sigma^{11} & \Sigma^{12} \\ \Sigma^{21} & \Sigma^{22} \end{bmatrix} \begin{bmatrix} \mathbf{x}_1 - \mu_1 \\ \mathbf{x}_2 - \mu_2 \end{bmatrix} \\
&= (\mathbf{x}_1 - \mu_1)^T \Sigma^{11} (\mathbf{x}_1 - \mu_1) + 2(\mathbf{x}_1 - \mu_1)^T \Sigma^{12} (\mathbf{x}_2 - \mu_2) + (\mathbf{x}_2 - \mu_2)^T \Sigma^{22} (\mathbf{x}_2 - \mu_2)
\end{aligned}$$

Here we have assumed

$$\Sigma^{-1} = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}^{-1} = \begin{bmatrix} \Sigma^{11} & \Sigma^{12} \\ \Sigma^{21} & \Sigma^{22} \end{bmatrix}$$

According to theorem 2, we have

$$\Sigma^{11} = (\Sigma_{11} - \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{12}^T)^{-1} = \Sigma_{11}^{-1} + \Sigma_{11}^{-1} \Sigma_{12} (\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1} \Sigma_{12}^T \Sigma_{11}^{-1}$$

$$\Sigma^{22} = (\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1} = \Sigma_{22}^{-1} + \Sigma_{22}^{-1} \Sigma_{12}^T (\Sigma_{11} - \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{12}^T)^{-1} \Sigma_{12} \Sigma_{22}^{-1}$$

$$\Sigma^{12} = -\Sigma_{11}^{-1} \Sigma_{12} (\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1} = (\Sigma^{21})^T$$

Substituting the second expression for Σ^{11} , first expression for Σ^{22} , and Σ^{12} into $Q(\mathbf{x}_1, \mathbf{x}_2)$ to get:

$$\begin{aligned}
Q(\mathbf{x}_1, \mathbf{x}_2) &= (\mathbf{x}_1 - \mu_1)^T [\Sigma_{11}^{-1} + \Sigma_{11}^{-1} \Sigma_{12} (\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1} \Sigma_{12}^T \Sigma_{11}^{-1}] (\mathbf{x}_1 - \mu_1) \\
&\quad - 2(\mathbf{x}_1 - \mu_1)^T [\Sigma_{11}^{-1} \Sigma_{12} (\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1}] (\mathbf{x}_2 - \mu_2) \\
&\quad + (\mathbf{x}_2 - \mu_2)^T [(\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1}] (\mathbf{x}_2 - \mu_2) \\
&= (\mathbf{x}_1 - \mu_1)^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1) \\
&\quad + (\mathbf{x}_1 - \mu_1)^T \Sigma_{11}^{-1} \Sigma_{12} (\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1} \Sigma_{12}^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1) \\
&\quad - 2(\mathbf{x}_1 - \mu_1)^T [\Sigma_{11}^{-1} \Sigma_{12} (\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1}] (\mathbf{x}_2 - \mu_2) \\
&\quad + (\mathbf{x}_2 - \mu_2)^T [(\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1}] (\mathbf{x}_2 - \mu_2) \\
&= (\mathbf{x}_1 - \mu_1)^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1) \\
&\quad + [(\mathbf{x}_2 - \mu_2) - \Sigma_{12}^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1)]^T (\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1} [(\mathbf{x}_2 - \mu_2) - \Sigma_{12}^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1)]
\end{aligned}$$

The last equal sign is due to the following equations for any vectors u and v and a symmetric matrix $A = A^T$:

$$\begin{aligned} u^T A u - 2u^T A v + v^T A v &= u^T A u - u^T A v - u^T A v + v^T A v \\ &= u^T A(u - v) - (u - v)^T A v = u^T A(u - v) - v^T A(u - v) \\ &= (u - v)^T A(u - v) = (v - u)^T A(v - u) \end{aligned}$$

We define

$$b \triangleq \mu_2 + \Sigma_{12}^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1)$$

$$A \triangleq \Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12}$$

and

$$\begin{cases} Q_1(\mathbf{x}_1) & \triangleq (\mathbf{x}_1 - \mu_1)^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1) \\ Q_2(\mathbf{x}_1, \mathbf{x}_2) & \triangleq [(\mathbf{x}_2 - \mu_2) - \Sigma_{12}^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1)]^T (\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12})^{-1} [(\mathbf{x}_2 - \mu_2) - \Sigma_{12}^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1)] \\ & = (\mathbf{x}_2 - b)^T A^{-1} (\mathbf{x}_2 - b) \end{cases}$$

and get

$$Q(\mathbf{x}_1, \mathbf{x}_2) = Q_1(\mathbf{x}_1) + Q_2(\mathbf{x}_1, \mathbf{x}_2)$$

Now the joint distribution can be written as:

$$\begin{aligned} f(\mathbf{x}) &= f(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{(2\pi)^{n/2} |\Sigma|^{1/2}} \exp\left[-\frac{1}{2} Q(\mathbf{x}_1, \mathbf{x}_2)\right] \\ &= \frac{1}{(2\pi)^{n/2} |\Sigma_{11}|^{1/2} |\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12}|^{1/2}} \exp\left[-\frac{1}{2} Q(\mathbf{x}_1, \mathbf{x}_2)\right] \\ &= \frac{1}{(2\pi)^{p/2} |\Sigma_{11}|^{1/2}} \exp\left[-\frac{1}{2} (\mathbf{x}_1 - \mu_1)^T \Sigma_{11}^{-1} (\mathbf{x}_1 - \mu_1)\right] \frac{1}{(2\pi)^{q/2} |A|^{1/2}} \exp\left[-\frac{1}{2} (\mathbf{x}_2 - b)^T A^{-1} (\mathbf{x}_2 - b)\right] \\ &= N(\mathbf{x}_1, \mu_1, \Sigma_{11}) N(\mathbf{x}_2, b, A) \end{aligned}$$

The third equal sign is due to theorem 3:

$$|\Sigma| = |\Sigma_{11}| |\Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12}|$$

The marginal distribution of \mathbf{x}_1 is

$$f_1(\mathbf{x}_1) = \int f(\mathbf{x}_1, \mathbf{x}_2) d\mathbf{x}_2 = \frac{1}{(2\pi)^{p/2} |\Sigma_{11}|^{1/2}} \exp\left[-\frac{1}{2}(\mathbf{x}_1 - \mu_1)^T \Sigma_{11}^{-1}(\mathbf{x}_1 - \mu_1)\right]$$

and the conditional distribution of \mathbf{x}_2 given \mathbf{x}_1 is

$$f_{2|1}(\mathbf{x}_2|\mathbf{x}_1) = \frac{f(\mathbf{x}_1, \mathbf{x}_2)}{f(\mathbf{x}_1)} = \frac{1}{(2\pi)^{q/2} |A|^{1/2}} \exp\left[-\frac{1}{2}(\mathbf{x}_2 - b)^T A^{-1}(\mathbf{x}_2 - b)\right]$$

with

$$b = \mu_2 + \Sigma_{12}^T \Sigma_{11}^{-1}(\mathbf{x}_1 - \mu_1)$$

$$A = \Sigma_{22} - \Sigma_{12}^T \Sigma_{11}^{-1} \Sigma_{12}$$

[next](#) [up](#) [previous](#)

Next: [Appendix B: Kernels and Up](#) **Up:** [Appendix A: Conditional and Previous:](#) [Inverse and determinant of](#)
Ruye Wang 2006-11-14